

Flavomycin® 4

(bambermycins
Type A medicated article)

ANTIBIOTIC

INDICATIONS FOR USE

Species	Bambermycins g/ton in Type C medicated feed	Indication(s)
Broiler Chickens	1-2*	Increased rate of weight gain; improved feed efficiency
Growing-Finishing Swine	2	Increased rate of weight gain; improved feed efficiency
	2-4*	Increased rate of weight gain
Growing Turkeys	2	Increased rate of weight gain; improved feed efficiency
	1-2*	Improved feed efficiency

*Only a single level of bambermycins (e.g., 4 grams/ton bambermycins) can be indicated on a feed tag. A different feed tag must be submitted for each level of bambermycins needed for your complete feeding program.

MIXING DIRECTIONS:

It is recommended that Flavomycin® 4 be diluted with a suitable grain carrier before addition to the final feed. A dilution of 1 part of Flavomycin® 4 and 9 parts grain carrier is the suggested working intermediate preblend. The table below shows Flavomycin® 4 and working intermediate preblend addition levels for various commonly used feeds:

Feed	Bambermycins in Type C Medicated Feed (g/ton)	Pounds Flavomycin® 4 per Ton Type C Medicated Feed	Pounds 1-9 Intermediate Preblend per Ton Type C Medicated Feed
Broiler Chickens	1	.25	2.50
	2	.50	5.00
Growing-Finishing Swine	2	.50	5.00
	3	.75	7.50
	4	1.00	10.00
Growing Turkeys	1	.25	2.50
	2	.50	5.00

Thoroughly mix both working intermediate preblend and finished Type C medicated feed to ensure complete and uniform distribution of the Flavomycin® 4.

FEEDING DIRECTIONS:
Type C medicated feed containing Flavomycin® 4 should be fed continuously as the sole ration.



Approved by FDA under NADA # 044-759

KCP-S00142
Rev. 03-2024

Flavomycin® 4

(bambermycins
Type A medicated article)

ANTIBIOTIC

INDICATIONS FOR USE

For increased rate of weight gain and improved feed efficiency in broiler chickens, growing-finishing swine and growing turkeys.

ACTIVE DRUG INGREDIENT

Bambermycins 4 grams per pound (8.8 grams per kilogram)

INGREDIENTS

Roughage products and/or calcium carbonate, dried bambermycins fermentation product, ethoxyquin (a preservative), mineral oil.

MIXING DIRECTIONS AND INDICATIONS ON REVERSE SIDE

WITHDRAWAL PERIODS
No withdrawal period is required when used according to labeling.

FOR USE IN MANUFACTURED FEEDS ONLY

USER SAFETY WARNINGS

Not For Human Use
Keep out of reach of children.

Restricted Drug (California) – Use only as directed

Net Weight – 50 pounds (22.68 kg)

Distributed by:
HUEPHARMA, INC.
Peachtree City, GA 30269

Bottom Print
Live Copy Area

Flavomycin® 4
(bambermycins
Type A medicated article)