

Tylovet[®] 100



Tylovet[®] 100

Tylosin Phosphate

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in Swine and Beef Cattle Feeds Only

Do Not Feed Undiluted

Equivalent to 100 g Tylosin per Pound

Swine:

For reduction in severity of effects of atrophic rhinitis.

For control of swine dysentery associated with *Brachyspira hyodysenteriae*.

For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with **Tylovet Soluble** (tylosin) drinking water.

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with **Tylovet Soluble** (tylosin) in drinking water.

Beef Cattle:

For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

Restricted Drug (California) - Use Only as Directed.

Approved by FDA under ANADA # 200-484

Huvepharma and Tylovet are registered trademarks of Huvepharma EOOD.

Distributed By:

Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc., at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Net Weight: 50 lbs (22.68 kg)



Tylovet[®] 100



Tylovet[®] 100

Type A Medicated Article



Tylovet[®] 100

Directions for Use

Tylovet[®] 100 Type A Medicated Article

Tylosin Phosphate

Read All Directions Carefully
Before Mixing and Feeding

Net Weight: **50 lbs**
(22.68 kg)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Active Drug Ingredient: Tylosin (as tylosin phosphate) . . . 100 g per lb

Ingredients: Roughage products, pregalatinized starch and dipotassium phosphate.

Caution: To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed. Do not use in a liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing excess of sodium metabisulfite.

Warning: Tylovet 100 may be irritating to unprotected skin and eyes. When mixing and handling Tylovet 100 use protective clothing, impervious gloves and a dust respirator. In case of accidental eye exposure, flush eyes with plenty of water. Exposed skin should be washed with plenty of soap and water. Remove and wash contaminated clothing. Seek medical attention if irritation becomes severe or persists. The safety data sheet (SDS) contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-877-994-4883.

For reduction in severity of effects of atrophic rhinitis. Feed 100 g of tylosin per ton (1.0 pound Tylovet 100 per ton) of complete feed. Feed continuously as the sole ration.

For control of swine dysentery. Feed 100 g of tylosin per ton (1.0 pound Tylovet 100 per ton) of complete feed for at least three weeks. Follow with 40 g tylosin per ton (0.4 pound Tylovet 100 per ton) of complete feed until pigs reach market weight.

For the treatment and control of swine dysentery. Feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of Tylovet 100) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as Tylovet Soluble) per gallon in drinking water for 3 to 10 days.

For control of porcine proliferative enteropathies (PPE, ileitis). Feed 100 g of tylosin per ton (1.0 pound Tylovet 100 per ton) of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton (1.0 pound Tylovet 100 per ton) of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Alternatively, feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of Tylovet 100) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as Tylovet Soluble) per gallon in drinking water for 3 to 10 days. Feed continuously as the sole ration when feeding Tylovet.

NOTE: Organisms vary in their degree of susceptibility to any chemotherapeutic. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reconfirmed.

Mixing and Feeding Directions for Beef Cattle Feeds	Tylovet 100 Per Ton Of Type C Feed	Tylosin Per Ton Of Type C Feed
For reduction in incidence of liver abscesses in beef cattle associated with <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium pyogenes</i> . Feed continuously as the sole ration.	0.08 to 0.10 lbs	8 to 10 g
Mixing Directions for Liquid Type B Cattle Feeds		
LIMITATIONS: 1. pH must be between 4.5 and 6.0. 2. For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute. Recirculate again in 10 minutes. 3. For liquid feeds stored in mechanical, air, or other agitation-type tank system: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used. 4. Liquid Type B Cattle Feeds must bear an expiration date of 31 days after the date of manufacture.		

NOT FOR HUMAN USE

Store at Room Temperature, 20 - 25 °C (68 - 77 °F). Avoid moisture.
Not to be used after the date printed on the bottom of the bag.
Rev. 06-2021



TAKE TIME
OBSERVE LABEL DIRECTIONS

Tylovet[®] 100

Tylosin Phosphate

Type A Medicated Article

LBb627v9Tyl100mp22.68-USA0621

Date	June 2021	Product	Tylovet 100 - 22.68 kg
Label identifier number	LBb627v9Tyl100mp22.68-USA0621	Country	USA
Dimensions	550x800x140 mm	Text size	
Cut:	900 mm	Font	
Join:	dx 100	Unroll - scheme	
Type of Packing	Bag	Logo	 HUVEPHARMA®
Colors	Pantone 313 C Pantone 355 C Black	Author	