

Monensin and Tylosin Medicated Cattle Feed - Coccidiosis

Type B Medicated Feed

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For Use in Cattle Feeds Only

Do Not Feed Undiluted

INDICATIONS

For prevention and control of coccidiosis due to *Eimeria bovis* and *E. zuernii* and reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Arcanobacterium (Actinomyces) pyogenes* in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS

| | |
|--------------------------------------|-------------------|
| Monensin, USP ¹ | 41 to 1600 g/ton* |
| Tylosin phosphate ² | 32 to 400 g/ton* |

GUARANTEED ANALYSIS

| | |
|--|---------|
| Crude Protein, not less than..... | % |
| Non-Protein Nitrogen (NPN) ³ , not more than..... | % |
| Crude Fat, not less than | % |
| Crude Fiber, not more than..... | % |
| Calcium, not less than..... | % |
| Calcium, not more than..... | % |
| Phosphorus, not less than..... | % |
| Salt ⁴ , not less than..... | % |
| Salt ⁴ , not more than..... | % |
| Sodium ⁵ , not less than | % |
| Sodium ⁵ , not more than | % |
| Potassium, not less than..... | % |
| Vitamin A ^{4,6} , not less than..... | I.U./lb |

³When added.

⁴If added.

⁵Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁶Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Mix 50 to 500 pounds of Type B medicated feed with 1950 to 1500 pounds of a non-medicated feed, respectively to yield a Type C feed with 10 to 40 grams per ton of monensin and 8 to 10 grams per ton of tylosin. It is recommended that Type B feeds containing more than 1440 g/ton be further diluted before mixing in the total mixed ration.

CAUTIONS

Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

WARNINGS

 A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. 

Approved by FDA under ANADA # 200-642

**MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345**

Net Weight lb (kg) on bag or bulk

Lot No. _____

*The medicated feed label must state a single drug concentration.

¹Sourced from Monovet®, ANADA # 200-639

²Sourced from Tylovet®, ANADA # 200-484

Monovet and Tylovet are registered trademarks of Huvepharma EOOD.