

# **Decoquinate and Monensin**

## **Type C Complete Medicated Cattle Feed**

### **INDICATIONS**

For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*, and improved feed efficiency in cattle being fed in confinement for slaughter.

### **ACTIVE DRUG INGREDIENTS<sup>1</sup>**

Decoquinate <sup>a</sup> .....	12.9 to 90.8 g/ton <sup>2</sup>
Monensin, USP <sup>b</sup> .....	5 to 30 g/ton <sup>2</sup>

### **GUARANTEED ANALYSIS**

Crude Protein, not less than .....	%
Crude Fat, not less than .....	%
Crude Fiber, not more than .....	%
Calcium, not less than .....	%
Calcium, not more than .....	%
Phosphorus, not less than .....	%
Potassium, not less than .....	%
Salt <sup>3</sup> , not less than .....	%
Salt <sup>3</sup> , not more than .....	%
Sodium <sup>4</sup> , not less than .....	%
Sodium <sup>4</sup> , not more than .....	%

<sup>3</sup>If added.

<sup>4</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

### **INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

### **FEEDING DIRECTIONS**

Feed only to cattle being fed in confinement for slaughter. Feed continuously as the sole ration to provide 22.7 mg decoquinate per 100 lb of body weight per day and 50 to 360 mg of monensin per head per day. Feed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The following table provides examples of feeding rates and amount of drug per ton of feed:

Animal Wt. (lbs.)	Feed Intake (% Body Wt)	Lbs. Feed Per Head /Day	Decoquinate (grams/ton)	Monensin, 100 mg/head/day (grams/ton)
500	1.5	7.5	30.3	26.7
	2.0	10.0	22.7	20.0
	3.0	15.0	15.1	13.3
	3.5	17.5	13.0	11.4
600	1.5	9.0	30.3	22.2
	2.0	12.0	22.7	16.7
	3.0	18.0	15.1	11.1
	3.5	21.0	13.0	9.5
700	1.5	10.5	30.3	19.0
	2.0	14.0	22.7	14.3
	3.0	21.0	15.1	9.5
	3.5	24.5	13.0	8.2
800	1.5	12.0	30.3	16.7
	2.0	16.0	22.7	12.5
	3.0	24.0	15.1	8.3
	3.5	28.0	13.0	7.1

## CAUTION

Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Monensin medicated cattle feed is safe only for use in cattle. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin may be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

## WARNINGS

Do not feed to animals producing milk for food. Do not feed to lactating dairy cattle.

Approved by FDA under ANADA # 200-652

**MANUFACTURED BY  
BLUE BIRD FEED MILL  
Any town, USA 12345**

Net Weight lb (kg) on bag or bulk

**Expiration Date:** 30 days after date of manufacture

Lot No. \_\_\_\_\_

<sup>1</sup>Actual label must bear the single concentration of each drug.

<sup>2</sup>Concentrations (grams per ton) must be such that feed delivers a total intake of 22.7 mg decoquinate per 100 pounds body weight per day and 50 to 360 mg monensin per head per day. Indicate in feeding directions amount of medicated feed to be consumed daily.

<sup>a</sup>Sourced from Deccox®, NADA # 039-417

<sup>b</sup>Sourced from Monovet®, ANADA # 200-639

Deccox is a registered trademark of Zoetis Inc.

Monovet is a registered trademark of Huvepharma EOOD.