

**This Bluebird Label Is Intended For Use Only With  
Monensin<sup>a</sup> and Ractopamine<sup>b</sup>**

**Liquid Heifer Supplement  
Medicated  
(Type C Medicated Feed)**

**For Beef and Dairy Heifers**

**INDICATIONS**

Heifers Fed in Confinement for Slaughter: For Increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

**ACTIVE DRUG INGREDIENTS**

Melengestrol acetate<sup>c</sup>.....(0.0000276-0.00022%)\*  
Each pound contains.....(0.125-1.0) mg. melengestrol acetate (select one)\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than .....	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	%
Crude Fat, not less than .....	%
Crude Fiber, not more than.....	%
Calcium, not less than.....	%
Calcium, not more than.....	%
Phosphorus, not less than.....	%
Salt <sup>2</sup> , not less than .....	%
Salt <sup>2</sup> , not more than.....	%
Sodium <sup>3</sup> , not less than .....	%
Sodium <sup>3</sup> , not more than.....	%
Potassium, not less than.....	%
Vitamin A <sup>2,4</sup> , not less than.....	I.U./lb
pH.....	4.0 to 8.0

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## **DIRECTIONS FOR USE**

Must be top dressed or mixed with a complete ration containing monensin (10 to 40 g/ton) and ractopamine (9.8 to 24.6 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 28 to 42 days on feed.

## **MIXING DIRECTIONS**

Mixing directions for liquid Type C feeds stored in recirculating tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as described in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

## **CAUTION**

Melengestrol Acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading.

Approved by FDA under ANADA # 200-659

## **MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Expiration date not required

Lot No. \_\_\_\_\_

\*Final printed label on formulated Type C medicated feed must bear a single concentration of each drug.

<sup>a</sup>Sourced from Monovet<sup>®</sup>, ANADA # 200-639

<sup>b</sup>Sourced from Actogain<sup>®</sup>, ANADA # 200-548

<sup>c</sup>Sourced from MGA<sup>®</sup>, NADA # 039-402

Monovet is a registered trademark of Huvepharma EOOD.

Actogain is a registered trademark of Zoetis Inc.

MGA is a registered trademark of Zoetis Inc.