

# **Monensin Medicated Feedlot Cattle Feed**

## **Type C Medicated Feed**

### **For Cattle Fed in Confinement for Slaughter (For prevention and control of coccidiosis)**

#### **INDICATIONS**

For prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*.

#### **ACTIVE DRUG INGREDIENT**

Monensin, USP<sup>a</sup> ..... 10 to 40 g/ton\*

#### **GUARANTEED ANALYSIS**

Crude Protein, not less than .....	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	%
Crude Fat, not less than .....	%
Crude Fiber, not more than .....	%
Calcium, not less than .....	%
Calcium, not more than .....	%
Phosphorus, not less than .....	%
Salt <sup>2</sup> , not less than .....	%
Salt <sup>2</sup> , not more than .....	%
Sodium <sup>3</sup> , not less than .....	%
Sodium <sup>3</sup> , not more than .....	%
Potassium, not less than .....	%
Vitamin A <sup>2,4</sup> , not less than .....	I.U./lb

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

#### **INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

#### **FEEDING DIRECTIONS**

Feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon the severity of challenge, up to maximum of 480 milligrams per head per day.

## **CAUTION**

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

## **WARNING**

 A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. 

Approved by FDA under ANADA # 200-662

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot No. \_\_\_\_\_

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

<sup>a</sup>Sourced from Monovet®, ANADA # 200-639

Monovet is a registered trademark of Huvepharma EOOD.