

**Ractopamine Medicated Finishing Cattle  
Feed – Top Dress  
(Weight Gain and Feed Efficiency)  
(ractopamine hydrochloride Type C top dress medicated feed)**

**INDICATIONS FOR USE**

For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter during the last 28 to 42 days on feed.

**ACTIVE DRUG INGREDIENT**

Ractopamine hydrochloride<sup>a</sup>.....not to exceed 800 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than.....	%
Crude Fat, not less than .....	%
Crude Fiber, not more than.....	%
Calcium, not less than.....	%
Calcium, not more than.....	%
Phosphorus, not less than.....	%
Salt <sup>2</sup> , not less than.....	%
Salt <sup>2</sup> , not more than.....	%
Sodium <sup>3</sup> , not less than.....	%
Sodium <sup>3</sup> , not more than .....	%
Potassium, not less than.....	%
Vitamin A <sup>2,4</sup> , not less than.....	I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed a minimum of 1.0 lb per head per day of Ractopamine Medicated Finishing Cattle Feed TD -Type C Top Dress Medicated Feed continuously to cattle fed in confinement for slaughter to provide 70 to 400 mg/head/day for the last 28 to 42 days on feed.

**CAUTION**

Not for animals intended for breeding.

## WARNINGS

### Withdrawal Periods

No withdrawal period is required when used according to labeling.

## USER SAFETY WARNINGS

The active ingredient in Optigrid® 45, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Optigrid 45 formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Optigrid 45, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The Safety Data Sheet (SDS) contains more detailed occupational safety information.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Huvepharma, Inc. at 1-877-994-4883.

Approved by FDA under ANADA # 200-679

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

### Net Weight lb (kg) on bag or bulk

Lot Number: (if applicable) \_\_\_\_\_  
Manufactured On:

\*The medicated feed label must state a single drug concentration.

<sup>a</sup> Sourced from Optigrid® 45, ANADA # 200-679

Optigrid is a registered trademark of Huvepharma EOOD.

Revision date: June 2020