

**This Bluebird Label Is Intended For Use Only With
Monensin^a and Ractopamine^b**

**Liquid Heifer Supplement
Medicated
(Type C Medicated Feed)**

For Beef and Dairy Heifers

INDICATIONS

Heifers Fed in Confinement for Slaughter: For Increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

ACTIVE DRUG INGREDIENTS

Melengestrol acetate^c.....(0.0000276-0.00022%)*
Each pound contains.....(0.125-1.0) mg. melengestrol acetate (select one)*

GUARANTEED ANALYSIS

Crude Protein, not less than	%
Non-Protein Nitrogen (NPN) ¹ , not more than	%
Crude Fat, not less than	%
Crude Fiber, not more than.....	%
Calcium, not less than.....	%
Calcium, not more than.....	%
Phosphorus, not less than.....	%
Salt ² , not less than	%
Salt ² , not more than.....	%
Sodium ³ , not less than	%
Sodium ³ , not more than	%
Potassium, not less than.....	%
Vitamin A ^{2,4} , not less than.....	I.U./lb
pH	4.0 to 8.0

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Must be top dressed or mixed with a complete ration containing monensin (10 to 40 g/ton) and ractopamine (9.8 to 24.6 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 28 to 42 days on feed.

MIXING DIRECTIONS

Mixing directions for liquid Type C feeds stored in recirculating tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as described in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

CAUTION

Melengestrol Acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading.

Approved by FDA under ANADA # 200-684

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Expiration date not required

Lot No. _____

*Final printed label on formulated Type C medicated feed must bear a single concentration of each drug.

^aSourced from Monovet[®], ANADA # 200-639

^bSourced from Optigrid[®], ANADA # 200-679

^cSourced from CycleGuard[®], ANADA # 200-399

Monovet, Optigrid and CycleGuard are registered trademarks of Huvepharma EOOD.