



1-877-994-4883

MAXTON, NC 28364

HUVEPHARMA, INC.

VLN/PCN 605/143150

5001101.03

LIVE OOCYSTS

COCCIDIOSIS VACCINE

10 x 10,000 DOSES/DOSIS

COCCIDIOSIS VACCINE

LIVE OOCYSTS

VACUNA PARA LA PREVENCIÓN DE LA COCCIDIOSIS AVIAR

OOCISTOS VIVOS



10 x 10,000 dose vials/10 mL

10 frascos con 10,000 dosis/10 mL

INDICATIONS:

For the vaccination of healthy turkeys one day of age or older against coccidiosis due to *E. adenoeides*, *E. meleagrimitis* and *E. gallopavonis*. This product license is conditional; potency has not been fully demonstrated. For more information regarding safety data, see productdata.aphis.usda.gov.

STORAGE INSTRUCTIONS: Store at 35-46°F (2-8°C). Do Not Freeze.

DOSAGE AND ROUTE OF ADMINISTRATION:

Administer by coarse spray at one day of age. For use in turkeys only. Shake vials well before using. Dilute vaccine with clean tap water using a clean container. Dye may be added as a marker. Dilute each 10,000 dose vial according to the spray equipment manufacturer's recommendations. When using a Huvepharma supplied spray cabinet, it is recommended to dilute 10,000 doses to 2,500 mL.

INGREDIENTS:

Coccidiosis vaccine contains live oocysts of *Eimeria adenoeides*, *Eimeria meleagrimitis* and *Eimeria gallopavonis*. Product contains gentamicin and amphotericin B.

PRECAUTIONS AND WARNINGS:

- Feed and water containing anti-coccidial drugs should not be given to turkeys receiving this vaccine for 21 days post-vaccination.
- Inactivate unused contents before disposal.
- Do not mix with other products, except as specified on this label.
- Use entire contents when first opened.
- In case of human exposure, contact a physician.
- Do not vaccinate within 21 days before slaughter.
- Federal regulations prohibit the repacking or sale of the contents of this carton in fractional units.
- Do not accept if the seal is broken.

ALWAYS READ, UNDERSTAND AND FOLLOW THE LABEL AND USE DIRECTIONS.

COCCIDIOSIS VACCINE

LIVE OOCYSTS